





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/00		(11) International Publication Number: WO 99/63975
A2		(43) International Publication Date: 16 December 1999 (16.12.99)
(21) International Application Number: PCT/EP99/04013 (22) International Filing Date: 10 June 1999 (10.06.99) (30) Priority Data: 98110709.7 10 June 1998 (10.06.98) EP 98113974.4 25 July 1998 (25.07.98) EP (71) Applicant (for all designated States except US): BIOGNOSTIK GESELLSCHAFT FÜR BIOMOLEKULARE DIAGNOS- TIK MBH [DE/DE]; Gerhard-Gerdes-Strasse 19, D-37079 Göttingen (DE). (72) Inventors; and (75) Inventors/Applicants (for US only): SCHLINGENSIEPEN, Karl-Hermann [DE/DE]; Pappelweg 3, D-37085 Göttingen (DE). SCHLINGENSIEPEN, Reimar [DE/DE]; Am Gold- graben 13, D-37073 Göttingen (DE). BRYSCH, Wolfgang [DE/DE]; Calsowstrasse 56, D-37085 Göttingen (DE). (74) Agent: MEYERS, Hans-Wilhelm; Postfach 10 22 41, D-50462 Köln (DE).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published Without international search report and to be republished upon receipt of that report.

(54) Title: A METHOD FOR STIMULATING THE IMMUNE SYSTEM



(57) Abstract

Medicament comprising a combination of at least one inhibitor of the effect of a substance negatively effecting an immune response, the substance selected from the group consisting of TGF-β and its receptors, VEGF and its receptors, interleukin 10 (IL-10) and its receptors, PGE₂ and its receptors, wherein the inhibitor has a molecular weight of less than 100 kDa and at least one stimulator positively effecting an immune response.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

A Method for Stimulating the immune system

Two different approaches have been used in the prior art to enhance the immune response against neoplastic cells. One approach uses the addition of cytokines like interleukin-2 (IL-2) or transfection of tumor cells and/or immune cells with genes coding for cytokines like IL-2 or other proteins enhancing the immune response like transfection of tumor cells with lymphotactin or like transfection of T-lymphocytes with CD-40 Ligand.

The second approach uses the inhibition of immunosuppressive molecules to enhance the body's immune response to tumor cells. Thus, J. NEUROSURG. 78 (1993) 944-51, Jachimczak et al. (1993) and WO 94/25588, Schlingensiepen et al. (1994) teach the use of antisense oligonucleotides targeted to TGF- β to reverse tumor-induced immunosuppression.

Several documents in the prior art teach that a combination of these two approaches is either not efficacious or is not beneficial over use of one of the two approaches used alone.

Thus, CANCER BIOTHER. 8(2), 1993, 159 - 170, Gridley et al., as well as CANCER BIOTHER. 9(4), 1994, 317-327, Mao et al., both teach that a combination of anti-transforming growth factor-beta antibody with IL-2 does not cause significant antitumor effects.

Furthermore, PROC. NATL. ACAD. SCI 93, (1996), 2909-2914, Fakhrai et al., teaches that a combination of transfection with genes encoding antisense sequences to transforming growth factor beta (TGF- β) TGF- β mRNA with transfection of IL-2 into tumor cells does not increase the immune response against the tumor compared to transfection with TGF- β antisense alone.

Surprisingly, in contrast, certain combinations of stimulators and inhibitors are more efficacious than either approach alone.

The present invention discloses a medicament comprising a combination of

- at least one inhibitor of the effect of a substance negatively effecting an immune response, the substance selected from the group consisting of TGF- β and its receptors, VEGF and its receptors, interleukin 10 (IL-10) and its receptors, PGE₂ and its receptors, wherein the inhibitor has a molecular weight of less than 100 kDa and
- at least one stimulator positively effecting an immune response.

In a preferred embodiment, the inhibitor is inhibiting the synthesis or function of molecules suppressing or downregulating or negatively affecting the immune response. The inhibitor can be an oligonucleotide which may function as an antisense nucleotide or a ribozyme or it may be an antibody fragment derived from an antibody e.g. a fab-fragment or a single chain antibody.

Preferably, the stimulator is positively effecting the immune response by increasing presentation of antigens and/or enhancing proliferation and/or function of immune cells.

In a preferred embodiment, the stimulator is enhancing the synthesis or function of molecules stimulating, enhancing, upregulating and/or positively regulating the immune response. In particular, the stimulator is stimulating and/or enhancing the synthesis and/or the function of factors such as GM-CSF, SCF, CSF, IFN- γ , FLT-3-ligand as well as monocyte chemotactic proteins (MCP-1), interleukin-2, interleukin-4, interleukin-12 and/or interleukin-18 or is one of the mentioned interleukins or is selected from the group consisting of viruses, viral antigens, antigens expressed in tumor cells or pathogens but not in normal cells, organspecific antigenes expressed in affected organs which are not essential for the organism, e. g. prostate, ovary, breast, melanine producing cells.

The stimulators are preferably selected from

- a) Chemokines, including lymphotactin and/or immune cell attracting substances and/or
- b) viruses and/or parts of viruses, including retroviruses, adenoviruses, papillomaviruses, Epstein-Barr-Viruses, Viruses that are non-pathogenic including Newcastle-Disease virus, Cow-pox-virus and/or
- c) autologous and/or heterologous MHC-Molecules and/or
- d) molecules involved in antigen processing and/or
- e) molecules involved in antigen presentation and/or
- f) molecules involved in mediating immune cell effects and/or
- g) molecules involved in mediating immune cell cytotoxic effects and/or

- h) molecules involved in antigen transportation and/or
- i) co-stimulatory molecules
- j) peptides enhancing recognition by immune cells and/or cytotoxic effects of immune cells
- k) the peptides containing one or more amino acids differing between a protein in the target cell from the other cells within an organism
- l) the peptides according to j) being
 - Peptides containing one or more mutations and/or amino acid substitutions of the ras protein amino and/or
 - Peptides containing one or more mutations and/or amino acid substitutions of the p53 protein and/or
 - Peptides containing one or more mutations and/or amino acid substitutions of the EGF-Receptor protein and/or
 - Peptides containing one or more mutations and/or amino acid substitutions of fusion peptides and/or fusion proteins and/or
 - Peptides containing one or more mutations and/or amino acid substitutions and/or amino acid substitutions caused by gene rearrangements and/or gene translocations and/or
 - Peptides containing one or more mutations and/or amino acid substitutions of the retinoblastoma protein and/or

- 5 -

- Peptides containing one or more mutations and/or amino acid substitutions of proteins coded by oncogenes and/or protooncogenes and/or
- Peptides containing one or more mutations and/or amino acid substitutions of proteins coded by anti-oncogenes and/or tumor suppressor genes and/or
- Peptides derived from proteins differing in the target cell by one or more amino acids from the proteins expressed by other cells in the same organism and/or
- Peptides derived from viral antigens and/or coded by viral nucleic acids and/or
- Peptides derived from proteins expressed in a diseased organ but not in the nervous system, muscle, hematopoietic system or other organs essential for survival. Diseased organs are e. g. prostate, ovary, breast, melanine producing cells and the like.

m) tumor cell extracts and/or tumor cell lysates and/or adjuvants;

n) fusion cells of dendritic and tumor cells.

These fusion cells are hybridoma cells derived from a mixture of dendritic cells and tumor cells. Dendritic cells are generated e. g. by treatment of PBMC with GM-CSF and IL-4 or a mixture of GM-CSF, IL-4 and IFN- γ or FLT-3 ligand. Fusion of dendritic cells with tumor cells can be achieved e. g. using PEG (polyethylene glycole) or electrofusion.

Surprisingly, treatment of PBMC with VEGF-oligonucleotides enhanced the number and/or effectiveness of dendritic cells.

In one embodiment of the invention the inhibitor is an oligonucleotide. Preferably the oligonucleotides of Fig. 1 are useful in the medicament of the present invention.

In a further embodiment, the invention provides oligonucleotides having one of the sequences given in figure 1-2 to 1-4.

Also oligonucleotides having 1 to 10 additional nucleotides at the 5'- or 3'- end are part of the invention.

Oligonucleotide sequences used for transfection are usually much longer sequences than those used for antisense oligonucleotides, which usually do not exceed 30 bases in length and are applied as short single-stranded sequences and are not integrated into a vector system.

Since transfected sequences are usually much longer than oligonucleotides, if cross inhibition of different members of a protein family would occur with the antisense technology, such cross inhibition of other mRNAs than the target mRNA, is much more likely with transfected antisense sequences, compared to oligonucleotides. However, Cell Growth Differ, Vol. 6(12), February 1995, pages 1635 - 1642, Huang, F. et al. teaches "only the K6 transfectant exhibited 39 and 33% respectively of the levels of TGF beta1 mRNA and active secreted TGF beta1 protein of the parental line. K6 exhibited no change in TGF beta2 expression and TGF beta3 expression was not detected in either parental or transfectant cell line."

It was therefore surprising to find oligonucleotides according to this invention, which were able to significantly reduce expression of both, TGF- β_1 as well as TGF- β_2 e. g. TGF- β_1 -14, TGF- β_1 -15, TGF- β -17-c-2260, TGF- β -123-2262, TGF- β -23-2268, TGF- β -2-4, TGF- β -2-14, TGF- β -2-15, TGF- β -2-9, TGF- β -2-14/1, TGF- β -2-14/2, TGF- β -1-136. Furthermore surprisingly oligonucleotides were designed, which were

able to significantly reduce expression of TGF- β_2 as well as TGF- β_3 .

Surprisingly even oligonucleotides were found, which were able to significantly reduce expression of TGF- β_2 as well as TGF- β_1 and TGF- β_3 , e. g. b1-N17, b1-N14, b1-N24, TGF- β_2 -9, TGF- β_2 -14, TGF- β_2 -15, TGF- β -17-c-2260, TGF- β -12-9/20-2261, TGF- β -123-2262, TGF- β -12-9/22-2263, TGF- β -23-2268, TGF- β_1 -98-11, TGF- β_1 -98-23, TGF- β_3 -98-7, TGF- β_3 -98-10, TGF- β -1-rwk-5, TGF- β -3-rwk-2, TGF- β -1-rwk-5, TGF- β -3-rwk-9, TGF- β -3-rwk-23, TGF- β_1 -3, TGF- β_1 -10.

Thus oligonucleotides which are effective against expression of at least two of TGF- β_1 , TGF- β_2 and/or TGF- β_3 are also part of the invention.

These findings were also surprising in view of the fact that sequence comparison between the mRNAs of TGF- β_2 , TGF- β_1 and TGF- β_3 showed that not a single sequence of 20 bases in length could be found that would be identical within the three different mRNAs. Even if such a hypothetical sequence had really existed, inhibition of the three mRNAs by such a hypothetical consensus sequence would have been extremely unlikely, since it is well known in the art that only a small minority of antisense sequences complementary to a certain mRNA actually exert a so-called antisense effect, *i.e.* inhibit expression of the respective protein.

Endothelial synthesis of monocyte chemotactic protein-1 (MCP-1) has been implicated in the regulation of monocyte recruitment for extravascular pools both under physiological and inflammatory conditions.

MCP-1 antisense oligonucleotides were able to modulate monocyte infiltration and were thus anti-inflammatory.

These antisense-oligonucleotides are useful for the treatment of inflammatory diseases e.g. asthma, morbus crohn, collitis ulcerosa, diabetes, glomerulonephritis, acute respiratory distress syndrome and arteriosclerotic plaque formation.

In a preferred embodiment of the invention the oligonucleotides and/or ribozymes and/or nucleic acids have modifications at the bases, the sugars and/or the phosphate moieties of the oligonucleotides.

In a further preferred embodiment of the invention the oligonucleotides and/or ribozymes and/or nucleic acids have modifications wherein the modifications are phosphorothioate (S-ODN) internucleotide linkages and/or methylphosphonate internucleotide linkages and/or phosphoramidate linkages and/or peptide linkages and/or 2'-O-derivatives, such as 2'-O-methyl or 2'-O-methoxyethoxy modifications of the sugar and/or modifications of the bases.

In a further preferred embodiment of the invention the oligonucleotides and/or ribozymes and/or nucleic acids are coupled to or mixed with folic acid, hormones, steroid hormones such as oestrogene, progesterone, corticosteroids, mineral corticoids, peptides, proteoglycans, glycolipids, phospholipids, polyethylene imine or other poly cations and derivatives therefrom.

Furthermore, the present invention provides a method of treating hyperproliferative diseases, neoplasms or infectious diseases by administering a medicament of the invention to patients in need thereof. The method is especially useful for the treatment of leukemia, non-hodgkin lymphoma, hodgkin lymphoma, bronchial carcinoma, esophageal carcinoma, colorectal carcinoma, gastric carcinomas, intestinal tumors, hepatic tumors, gall bladder and gallduct carcinomas, pancreatic carcinoma, anal carcinoma, breast cancer, ovarian carcinoma, cervical carcinoma, endometrium carcinoma, prostatic carcinoma, bladder carcinoma, malignant melanoma, brain tumors, and sarcomas.

The necessary doses of the medicament of the present invention depend on the disease and the severity of the disease. Whereas higher levels are more effective, they often have a higher degree of side effects. Suitable doses are selected to obtain concentrations of the oligonucleotides in the range of 0.1 to 10 $\mu\text{mol/l}$ and concentrations of the cytokines in the range of 10 to 1.000 U/ml in the patient blood.

In a preferred embodiment of the invention the inhibitor of the effect of a substance negatively effecting an immune response is applied locally to a tumor or other pathologically affected site or organ and the stimulator positively effecting an immune response is applied systemically (e.g. i.v. or s.c. or orally).

In another preferred embodiment of the invention the inhibitor of the effect of a substance negatively effecting an immune response is applied systemically (e.g. i.v. or s.c. or orally) to the tumor and the stimulator positively effecting an immune response is applied locally to a tumor or other pathologically affected site or organ.

In another preferred embodiment of the invention the inhibitor of the effect of a substance negatively effecting an immune response is applied systemically (e.g. i.v. or s.c. or orally) to the tumor and the stimulator positively effecting an immune response is applied systemically (e.g. i.v. or s.c. or orally).

In another preferred embodiment of the invention the inhibitor of the effect of a substance negatively effecting an immune response is applied locally to a tumor or other pathologically affected site or organ and the stimulator positively effecting an immune response is applied locally to a tumor or other pathologically affected site or organ.

Fig. 1 shows oligonucleotides useful in the present invention.

Fig. 2A shows effects of oligonucleotides (f.c. 5 μ M) on TGF- β 2 secretion in glioma cells in 10% MEM Dulbecco medium (3 day incubation with oligonucleotides).

Fig. 2B shows effects of oligonucleotides (f.c. 5 μ M) on TGF- β 1 secretion in PBMC in 10% FCS RPMI 1640 medium (3 day incubation with oligonucleotides).

Fig. 3A shows effects of oligonucleotides (f.c. 5 μ M) on TGF- β 1 secretion in PBMC in 10% FCS RPMI 1640 medium (3 day incubation with oligonucleotides).

Fig. 3B shows effects of oligonucleotides (f.c. 5 μ M) on TGF- β 2 secretion in glioma cells in 10% FCS RPMI 1640 medium (3 day incubation with oligonucleotides).

Fig. 4A shows TGF- β 1 concentration (ELISA) in glioma cells (3 day incubation with oligonucleotides).

Fig. 4B shows TGF- β 2 concentration (ELISA) in glioma cells (3 day incubation with oligonucleotides).

Fig. 5 shows lysis of tumor-cells: LAK-Cytotoxicity, Ratio of glioma-cells/PBMC: 1 : 20.

Fig 6A shows dendritic cells generated from PBMC (% of control). Cytokines: GM-CSF (400 U/ml) + IL-4 (300 U/ml).

Fig. 6B shows lysis of tumor-cells: Effects of 5 μ M VEGF-Antisense-Oligos on LAK-Cytotoxicity. Ration of tumor-cells/DC/PBMC was 1 : 5 :20.

Fig. 7A shows effects of oligonucleotides (f.c. 5 μ M) on TGF- β 1 secretion in PBMC in 10% FCS RPMI 1640 medium (3 day incubation with oligonucleotides).

Fig. 7B shows effects of oligonucleotides (f.c. 5 μ M) on TGF- β 2 secretion in tumor cells in 10% FCS RPMI 1640 medium (3 day incubation with oligonucleotides).

Fig. 8 shows lysis of tumor-cells: Effects of oligonucleotides on LAK-Cytotoxicity. Ration of tumor-cells/PBMC was 1 : 20.

Examples

Preparation of PBMC and tumor cells

Peripheral blood mononuclear cells (PBMC) were isolated from venous blood of healthy donors by standard Ficoll-Hypaque gradient centrifugation. Briefly, heparinized blood was mixed with equal volumes of complete medium (CM: RPMI 1640 medium supplemented with 10% (v/v) fetal calf serum and 1 mM L-Glutamine) and layered onto a Ficoll-Hypaque (Pharmacia, Uppsala, Sweden) gradient. After centrifugation at 400g for 30 min at room temperature, PBMCs banded at the plasma-Ficoll interface were recovered, washed three times and resuspended in complete medium. Cell viability, as determined by Trypan blue exclusion, was greater than 97%.

Human glioma cell lines were established from tumor specimens of patients with anaplastic astrocytoma (WHO Grad III) or from glioblastoma (WHO Grad IV).

Measurement of cell proliferation

For PBMC-proliferation assays (3H-thymidine incorporation and cell counting), freshly isolated PBMCs were cultured for 72h in 96-well round-bottom plates (Nunc, Copenhagen, Denmark) at a final concentration (f.c.) of 10^5 cells/well (100 μ l CM). For cell number determination the cells were counted by hemacytometer. Cell viability was determined by trypan blue staining. Treated and untreated cells showed 95-100% viability after 72h *in vitro* growth (with or without S-ODN).

For the tumor proliferation experiments $10^4/100$ μ L glioma cells were seeded into 96-well flat-bottom plates (Nunc, Denmark) and incubated with cytokines and/or oligonucleotides. The DNA synthesis rate was measured by a standard 3H-thymidine incorporation assay and determination of cell number was performed as described above.

Quantification of TGF- β 1 protein in culture supernatants by enzyme-linked immunosorbent assay (ELISA)

The culture medium was harvested after 3 days, cleared of cellular components by centrifugation, filtered and stored at -70°C until processed further. TGF- β 1 and TGF- β 2 concentrations were measured after acidification of supernatants by TGF- β 1 and TGF- β 2 ELISA (R&D Systems, Minneapolis, USA) in duplicates, as recommended by the manufacturer.

Figures 1 - 4 and 7 show the effect of oligonucleotides on the TGF- β secretion in cells. The concentration of the TGF- β is reported as an optical density. The higher the optical density the higher is the concentration of the TGF- β .

Figure 1A and 1B shows the effect of the oligonucleotides on the TGF- β secretion. Control oligos (GAA GGA ATT ACC ACT TTC) have no effects whereas the oli-

oligonucleotides shown in the figures reduce the secretion of TGF- β . The oligos in figure 1 are more effective against TGF- β 1.

Figure 2 shows further oligos and their effects on TGF- β secretion. TGF- β -14 is especially effective against the secretion of TGF- β 1 and - β 2.

Figure 3 shows further oligonucleotides being effective against secretion of TGF- β 1 and - β 2. These oligonucleotides are more effective against TGF- β 2 but are also effective against TGF- β 1.

Figure 8 shows a supra additive effect on tumor cell cytotoxicity by a combination of 2 μ M each of a TGF- β 1 and TGF- β 2 antisense oligonucleotide compared to a single 5 μ M dose of either oligonucleotide.

CARE-LASS (calcein-release-assay) to measure cytotoxic PBMC activity

A standard calcein-release-assay (CARE-LASS assay) to determine cytotoxic activity of PBMC was employed as described by Lichtenfels, R., Biddison, W.E., Schulz, H., Vogt, A.B. and R. Martin. CARE-LASS (calcein-release assay), an improved fluorescence-based test system to measure cytotoxic lymphocyte activity. J. Immunol. Meth., 172: 227-239, 1994.

Target and Effector cells

At the day of the assay malignant glioma were harvested, washed twice in 5% FCS /PBS and incubated with Calcein-AM (Molecular Probes, USA) for 30min in 37°C. Labeled target cells were washed twice in 5% FCS/PBS, adjusted to 100 000 / ml, and plated into 96-well U-shaped microtiter plates (Nunc, Denmark) at the final volumen of 100uL/well.

PBMC were washed with 5% FCS/PBS and adjusted to final concentration of 1-10 Mio cells /ml.

Cells were treated with cytokines and oligodeoxynucleotides as described in the individual experiments.

Assay

To measure CTL activity effector cells were plated into 96-well U-shape microtiter plates at Target : Effector Ratios of 1:10 - 1:100. To measure spontaneous release and total release of calcein, wells were preloaded with 200uL 5%FCS/PBS or 200uL lysis buffer (50mM sodium-borate, 0.1% Triton, pH 9.0) respectively. After incubating the plate for 4 h at 37°C in an incubator, 100uL of supernatans were transferred into new wells and measured employing an automated fluorescence scanner (Titertek Fluoroskan II, Germany). Both for excitation and for emission, filter settings 2 were chosen (ex 2 - 485nm, em 2 -538 nm). The percent of cytotoxicity was determined from the following equation:

$$\frac{\text{F/CTL assay} - \text{F spontaneous release}}{\text{F total lysis} - \text{F spontaneous release}} \times 100 = \% \text{ cytotoxicity}$$

In one set of experiments, glioma cells, denritic cells (DC) and PBMC were co-cultured. In these experiments DC were generated from PBMC using the cytokines GM-CSF and IL4. Cells were further treated with antisense VEGF-oligonucleotides according to the invention or with no oligonucleotides as control experiments. Tumor cells were also treated with the cytokines GM-CSF and IL4 with or without oligonudeotides.

PBMC were only treated with oligonucleotides according to the invention, but not with the cytokines GM-CSF and IL4. oligos were used at a concentration of 5 μ M unless indicated otherwise in the descriptions in the figures.

The CARE-LASS (calcein-release-assay) was used to measure cytotoxic PBMC activity.

In one set of experiments glioma cells and PBMC were treated either with a single oligonucleotide or with a combination of oligonucleotides. The single oligonucleotides were given at 5 μ M concentration. In the combination experiment, each oligonucleotide was given at 2 μ M concentration. Both, PBMC and tumor cells were incubated separately with the oligonucleotide(s) for 72 h.

The CARE-LASS (calcein-release-assay) was used to measure cytotoxic PBMC activity.

Claims

1. Medicament comprising a combination of
 - at least one inhibitor of the effect of a substance negatively effecting an immune response, the substance selected from the group consisting of TGF- β and its receptors, VEGF and its receptors, interleukin 10 (IL-10) and its receptors, PGE₂ and its receptors, wherein the inhibitor has a molecular weight of less than 100 kDa and
 - at least one stimulator positively effecting an immune response.
2. The medicament of claim 1 wherein the inhibitor is inhibiting the synthesis or function of molecules suppressing or downregulating or negatively affecting the immune response.
3. The medicament of claim 1 wherein the inhibitor is an oligonucleotide.
4. The medicament according to claim 3 wherein the oligonucleotide is an antisense nucleotide and/or a ribozyme.
5. The medicament according to claim 3 wherein the oligonucleotides has a sequence according to figure 1.
6. The medicament according to claim 1, wherein the inhibitor is a fab-fragment or single chain antibody (scFv).
7. The medicament according to claim 1, wherein the stimulator is enhancing the synthesis or function of molecules stimulating, enhancing, upregulating and/or positively regulating the immune response.

8. The medicament according to claim 7, wherein the stimulator is stimulating and/or enhancing the synthesis and/or the function of factors such as GM-CSF, SCF, CSF, IFN, FLT-3-ligand, monocyte chemotatic proteins (MCP-1), interleukin-2, interleukin-4, interleukin-12 and/or interleukin-18 or is one of the mentioned interleukins or is selected from the groups consisting of viruses, viral antigens, antigens expressed in tumor cells or pathogens, but not in normal cells, organ specific antigens expressed in affected organs which are not essential for the organism or fusion cell of dendritic and tumor cells.
9. The medicament according to claim 1, wherein the medicament comprises two or more of the inhibitors and/or the stimulators.
10. An oligonucleotide having one of the sequences of figures 1-2 to 1-4 (No. 55 - 213).
11. The oligonucleotide according to claim 10 wherein each oligonucleotide is effective against expression of at least two of TGF- β_1 , TGF- β_2 and/or TGF- β_3 having the sequence.
12. A method of treating neoplasm or infectious diseases by administering a medicament according to claim 1 to a patient in need thereof.
13. A method according to claim 12 for the treatment of hyperproliferative diseases, leukemia, non-hodgkin lymphoma, hodgkin lymphoma, bronchial carcinoma, esophageal carcinoma, colorectal carcinoma, gastric carcinomas, intestinal tumors, hepatic tumors, gall bladder and gallduct carcinomas, pancreatic carcinoma, anal carcinoma, mastocarcinoma, ovarian carcinoma, cervical carcinoma, endometrium carcinoma, prostatic carcinoma, bladder carcinoma, malignant melanoma, brain tumors, and/or sarcomas.

1.	TGF-β2-1	C ACA CAG TAG TGC A
2.	TGF-β2-2	GC ACA CAG TAG TGC
3.	TGF-β2-3	GC TTG CTC AGG ATC TGC
4.	TGF-β2-4	TAC TCT TCG TCG CT
5.	TGF-β2-5	C TTG GCG TAG TAC T
6.	TGF-β2-6	G TAA ACC TCC TTG G
7.	TGF-β2-7	GT CTA TTT TGT AAA CCT CC
8.	TGF-β2-8	GC ATG TCT ATT TTG TAA ACC
9.	TGF-β2-9	CGG CAT GTC TAT TTT GTA
10.	TGF-β2-10	G GCA TCA AGG TAC C
11.	TGF-β2-11	CTG TAG AAA GTG GG
12.	TGF-β2-12	AC AAT TCT GAA GTA GGG T
13.	TGF-β2-13	T CAC CAA ATT GGA AGC AT
14.	TGF-β2-14	GCT TTC ACC AAA TTG GAA GC
15.	TGF-β2-15	CTG GCT TTT GGG TT
16.	TGF-β2-16	T CTG ATA TAG CTC AAT CC
17.	TGF-β2-17	T CCT AGT GGA CTT TAT AG
18.	TGF-β2-18	T TTT TCC TAG TGG ACT
19.	TGF-β2-19	C AAT TAT CCT GCA CAT TTC
20.	TGF-β2-20	GC AAT TAT CCT GCA CA
21.	TGF-β2-21	GC AGC AAT TAT CCT GC
22.	TGF-β2-22	TG GCA TTG TAC CCT
23.	TGF-β2-23	TG TGC TGA GTG TCT
24.	TGF-β2-24	CC TGC TGT GCT GAG TG
25.	TGF-β2-25	C TTG GGT GTT TTG C
26.	TGF-β2-26	T TTA GCT GCA TTT GCA AG
27.	TGF-β2-27	G CCA CTT TTC CAA G
28.	TGF-β2-14/1	CTT TCA CCA AAT TGG AAG
29.	TGF-β2-14/2	CAC CAA ATT GGA AGC
30.	TGF-β2-14/3	TCA CCA AAT TGG AAG C
31.	TGF-β2-15/1	CTC TGG CTT TTG GG
32.	TGF-β2-9/1	CGG CAT GTC TAT TTT G
33.	TGF-β1-1	CGA TAG TCT TGC AG
34.	TGF-β1-2	GTC GAT AGT CTT GC
35.	TGF-β1-3	CTT GGA CAG GAT CT
36.	TGF-β1-4	CCA GGA ATT GTT GC
37.	TGF-β1-5	CCT CAA TTT CCC CT
38.	TGF-β1-6	GAT GTC CAC TTG CA
39.	TGF-β1-7	CTC CAA ATG TAG GG
40.	TGF-β1-8	ACC TTG CTG TAC TG
41.	TGF-β1-9	GTA GTA CAC GAT GG
42.	TGF-β1-10	CAC GTA GTA CAC GA
43.	TGF-β1-11	CAT GTT GGA CAG CT
44.	TGF-β1-12	GCA CGA TCA TGT TG
45.	TGF-β1-13	TGT ACT CTG CTT GAA C
46.	TGF-β1-14	CTG ATG TGT TGA AGA ACA
47.	TGF-β1-15	CTC TGA TGT GTT GAA G
48.	TGF-β1-16	GGA AGT CAA TGT ACA G
49.	TGF-β1-17	CAT GTC GAT AGT CTT GCA
50.	TGF-β1-18	AGC TGA AGC AAT AGT TGG
51.	TGF-β1-19	GTC ATA GAT TTC GTT GTG
52.	TGF-β1-20	CTC CAC TTT TAA CTT GAG
53.	TGF-β1-21	TGC TGT ATT TCT GGT ACA
54.	TGF-β1-137	CGA TAG TCT TGC AG

55.	b1-N17	TCC TCT TCG ACT GCT CTC
56.	b2-N14	CGA AGG TTA AAC CAC TTT CG
57.	b2-N24	GTG AGT CGT GTC GTC C
58.	TGF-β2-98-1	CATCGTTGTCGTCG
59.	TGF-β2-98-2	CGCTTCTCCGCCG
60.	TGF-β2-98-3	CGAAGGAGAGCCATTCG
61.	TGF-β2-98-4	CGATGTAGCG
62.	TGF-β2-98-5	CGTCAAATCG
63.	TGF-β2-98-6	CGTAGTACTCTTCGTCG
64.	TGF-β2-98-7	CGCGCTCGCAGGCG
65.	TGF-β2-98-8	CGGCCGCCCTCCGGCTCG
66.	TGF-β2-98-9	CGCGGATCGCCTCG
67.	TGF-β2-98-10	GAGCGCGACCGTGAC
68.	TGF-β-17-c-2260	ACC TCC TTG GCG TAG TA
69.	TGF-β-12-9/20-2261	AGG GCG GCA TGT CTA TTT TG
70.	TGF-β-123-2262	CAG AAG TTG GCA TTG TAC
71.	TGF-β-12-9/22-2263	AGG GCG GCA TGT CTA TTT TGT A
72.	TGF-β-23-2268	TGG GAC ACG CAG CAA GG
73.	TGF-β1-98-1	CGGGGGCGGGGCGGGG
74.	TGF-β1-98-2	CGGGGGCGGGGCGGGGCG
75.	TGF-β1-98-3	CGGCGCCGCCGAGGCGCCCCG
76.	TGF-β1-98-4	CCGAGGTCTTGC GG
77.	TGF-β1-98-5	CGGCGGTGCCGGGA
78.	TGF-β1-98-6	CTCGGCGGCCGGTAG
79.	TGF-β1-98-7	CGCTAAGGCG
80.	TGF-β1-98-8	CCGCACA ACTCCGG
81.	TGF-β1-98-9	GCGAGTCGCTGG
82.	TGF-β1-98-10	CGGTTGCTGAGGTATCG
83.	TGF-β1-98-11	CCGGGAGAGCAACACGG
84.	TGF-β1-98-12	CGCTTCTCG
85.	TGF-β1-98-13	CCATTAGCACGCGGG
86.	TGF-β1-98-14	CGGGCTCCG
87.	TGF-β1-98-15	CCGGCCACCCGGTCGCGG
88.	TGF-β1-98-16	CGAGCACGGCCTCG
89.	TGF-β1-98-17	CGGGCAGCGGGCCGGGCG
90.	TGF-β1-98-18	CGCGGATGGCCTCG
91.	TGF-β1-98-19	CGATGCGCTTCCG
92.	TGF-β1-98-20	CCCGCGGCCGGCGGG
93.	TGF-β1-98-21	CGCAGCCCGGAGGGCG
94.	TGF-β1-98-22	CGGCGCCCCCG
95.	TGF-β1-98-23	CGGCACTGCCGAGAGCGCG
96.	TGF-β1-98-24	CGGGGATGAAGGCGGCG
97.	TGF-β1-98-25	CGGGTCGGCGACTCCCG
98.	TGF-β1-98-26	CGCCTGAGGGACGCCG
99.	TGF-β1-98-27	AAGCGTCCCCGGCG
100.	TGF-β1-98-28	CGCGGGGCAGCGTCGCG
101.	TGF-β1-98-29	CCCCGCGCCTCCGG
102.	TGF-β1-98-30	CGGCGGGCGGCTCG
103.	TGF-β1-98-31	CGCTCCGGGCCGAGGCCG
104.	TGF-β1-98-32	CGGCCCCGCGGGCG
105.	TGF-β1-98-33	CGGACGGGGCGTCC
106.	TGF-β1-98-34	CGGCCGGGGCCCTCG

Fig. 1-2

107.	TGF-β3-98-1	TCGAGCTTCCCCGA
108.	TGF-β3-98-2	CCCGGAGCCGAAGG
109.	TGF-β3-98-3	CCCGAGGAGCGGG
110.	TGF-β3-98-4	ACGCAGCAAGGCGA
111.	TGF-β3-98-5	CGGGTTGTGAGCCG
112.	TGF-β3-98-6	CGGCAGTGCCCCG
113.	TGF-β3-98-7	CGGAATTCTGCTCG
114.	TGF-β3-98-8	TTCGTTGTGCTCCG
115.	TGF-β3-98-9	ATTCCGACTCGGTG
116.	TGF-β3-98-10	ACGTGGGTCATCACCGT
117.	TGF-β3-98-11	CGAAGAAGCG
118.	TGF-β3-312	CCT AAT GGC TTC CA
119.	VEGF-98-1	CGGCCGCGGTGTGT
120.	VEGF-98-2	CGGGAATGCTTCCGCCG
121.	VEGF-98-3	CGGCTCACCGCCTCGGC
122.	VEGF-98-4	CACGTCTGCGGATC
123.	VEGF-98-5	CCCCGCATCGCATCAGGG
124.	VEGF-98-6	CGCCTTGCAACGCG
125.	VEGF-98-7	CCGACCGGGGCCCG
126.	VEGF-49	GTTTCATGGTTTCGG
127.	VEGF-55	GCAGAAAGTTCATGG
128.	VEGF-188	GCTGATAGACATCC
129.	VEGF-190	GCGCTGATAGACAT
130.	VEGF-194	GTAGCTGCGCTGATAG
131.	VEGF-253	CTCGATCTCATCAG
132.	VEGF-255	ATGTACTCGATCTCATC
133.	VEGF-260	GAAGATGTACTCGATC
134.	VEGF-263	CTTGAAGATGTACTCG
135.	VEGF-292	GCATCGCATCAGGG
136.	VEGF-294	CCGCATCGCATCAG
137.	VEGF-422	CATTTGTTGTGCTGTAGG
138.	VEGF-434	GGTCTGCATTCACATTTG
139.	VEGF-441	CTTTGGTCTGCATTC
140.	VEGF-445	CTTTCTTTGGTCTGC
141.	VEGF-450	GCTCTATCTTTCTTTGG
142.	VEGF-455	GTCTTGCTCTATCTTTC
143.	VEGF-459	CTTGTCTTGCTCTATC
144.	VEGF-596	CATCTGCAAGTACGTTTCG
145.	VEGF-598	CACATCTGCAAGTACGTT
146.	VEGF-600	GTCACATCTGCAAGTACG
147.	VEGF-600-2	CATCTGCAAGTACG
148.	VEGF-601	CACATCTGCAAGTAC
149.	VEGF-604	GTCACATCTGCAAG
150.	VEGF-607	CTTGTACATCTGC
151.	VEGF-607-2	GGCTTGTCACATCTGC
152.	VEGF-610	CTCGGCTTGTCACATC
153.	VEGF-638	CTCCTTCCTCCTGC
154.	VEGF-766	GCT TGA AGA TGT ACCT CG
155.	VEGF-r-1062	CGT TGC TCT CCG ACG
156.	flt-1165	GAC ACG GCC TTT TCG
157.	flt-rm-2115	CCA GCA GCT GAC CAT GG
158.	flkl1/kdr-m-2315	GAA ATC GAC CCT CGG
159.	MCP-1-Rec-A/B-571	GCA TGT TGT GGA TG
160.	MCP-1-1954	GCA GAG ACT TTC ATG C
161.	MCP-1-1955	ATA ACA GCA GGT GAC TGG

Fig. 1-3

162.	MCP-1-1956	GAA CCC ACT TCT GC
163.	MCP-1-2761	GAC ACT TGC TGC TG
164.	MCP-1-2762	CCA CTT CTG CTT GGG
165.	VEGF-703	CTG CAA GTA CGT TCG
166.	flt-1567	TCC CTT ATG ATG CCA GCA AGT, G
167.	TGF- β -Rec-I-796	CCA GCA ATG ACA GC
168.	TGF- β -1-rwk-1	G GGA AAG CTG AGG C
169.	TGF- β -1-rwk-2	T CGA GGG AAA GCT GA
170.	TGF- β -1-rwk-3	C CTC GAG GGA AAG C
171.	TGF- β -1-rwk-4	GG GCT GGT GTG GTG
172.	TGF- β -1-rwk-5	GA ACA GGG CTG GTG TG
173.	TGF- β -1-rwk-6	G AAC AGG GCT GGT G
174.	TGF- β -1-rwk-7	AG AGC GCG AAC AGG
175.	TGF- β -1-rwk-8	GA GAG CGC GAA CAG G
176.	TGF- β -1-rwk-9	CGA GAG CGC GAA CAG
177.	TGF- β -1-rwk-10	CCC CTG GCT CGG GGG
178.	TGF- β -1-rwk-11	C CCT GGC TCG GGG
179.	TGF- β -1-rwk-12	C CCC TGG CTC GGG G
180.	TGF- β -1-rwk-13	TCC CCC TGG CTC GG
181.	TGF- β -1-rwk-14	C TCC CCC TGG CTC G
182.	TGF- β -1-rwk-15	TGC GCT TCC GCT TCA C
183.	TGF- β -1-rwk-16	CC TCG ATG CGC TTC
184.	TGF- β -1-rwk-17	G ATG GCC TCG ATG C
185.	TGF- β -1-rwk-18	G GAT GGC CTC GAT GC
186.	TGF- β -1-rwk-19	ATG GCC TCG ATG CGC TT
187.	TGF- β -3-rwk-1	TC AGC AGG GCC AGG
188.	TGF- β -3-rwk-2	GCA AAG TTC AGC AGG GC
189.	TGF- β -3-rwk-3	GG CAA AGT TCA GCA GG
190.	TGF- β -3-rwk-4	GT GGC AAA GTT CAG CAG G
191.	TGF- β -3-rwk-5	GTG GCA AAG TTC AG
192.	TGF- β -3-rwk-6	GAC CGT GGC AAA GTT CAG
193.	TGF- β -3-rwk-7	AGA GAG GCT GAC CGT
194.	TGF- β -3-rwk-8	GAC AGA GAG AGG CTG AC
195.	TGF- β -3-rwk-9	A CAG AGA GAG GCT GA
196.	TGF- β -3-rwk-10	GT GGA CAG AGA GAG G
197.	TGF- β -3-rwk-11	CA AGT GGA CAG AGA GAG G
198.	TGF- β -3-rwk-12	TCT TCT TGA TGT GGC C
199.	TGF- β -3-rwk-13	CC CTC TTC TTC TTG ATG
200.	TGF- β -3-rwk-14	C ACC CTC TTC TTC T
201.	TGF- β -3-rwk-15	A TGG ATT TCT TTG GCA T
202.	TGF- β -3-rwk-16	GGA TTT CTT TGG C
203.	TGF- β -3-rwk-17	AA GTT GGA CTC TCT TCT C
204.	TGF- β -3-rwk-18	TAA GTT GGA CTC TCT TCT
205.	TGF- β -3-rwk-19	GAC CTA AGT TGG ACT C
206.	TGF- β -3-rwk-20	T TTC TAG ACC TAA GTT GG
207.	TGF- β -3-rwk-21	CT GAT TTC TAG ACC TAA G
208.	TGF- β -3-rwk-22	G AAG CAG TAA TTG GTG T
209.	TGF- β -3-rwk-23	GG AAT CAT CAT GAG G
210.	TGF- β -3-rwk-24	GGG AAT CAT CAT GAG
211.	TGF- β -3-rwk-25	G GTT GTC GAG CCG GT
212.	TGF- β -3-rwk-26	GTC CTC CCA ACA TAG TA
213.	TGF- β -3-rwk-27	GG GTC CTC CCA ACA




Figure 2




Figure 3




Figure 4





Figure 5

B.



A.




Figure 6




Figure 7






Figure 8



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 48/00, 31/70, 38/20, 38/19, C12N 15/11 // (A61K 38/20, 31:70) (A61K 38/19, 31:70)		A3	(11) International Publication Number: WO 99/63975
			(43) International Publication Date: 16 December 1999 (16.12.99)
(21) International Application Number: PCT/EP99/04013		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 10 June 1999 (10.06.99)		Published With international search report.	
(30) Priority Data: 98110709.7 10 June 1998 (10.06.98) EP 98113974.4 25 July 1998 (25.07.98) EP		(88) Date of publication of the international search report: 3 August 2000 (03.08.00)	
(71) Applicant (for all designated States except US): BIOGNOSTIK GESELLSCHAFT FÜR BIOMOLEKULARE DIAGNOSTIK MBH [DE/DE]; Gerhard-Gerdes-Strasse 19, D-37079 Göttingen (DE).			
(72) Inventors; and (75) Inventors/Applicants (for US only): SCHLINGENSIEPEN, Karl-Hermann [DE/DE]; Pappelweg 3, D-37085 Göttingen (DE). SCHLINGENSIEPEN, Reimar [DE/DE]; Am Goldgraben 13, D-37073 Göttingen (DE). BRYSCH, Wolfgang [DE/DE]; Calsowstrasse 56, D-37085 Göttingen (DE).			
(74) Agent: MEYERS, Hans-Wilhelm; Postfach 10 22 41, D-50462 Köln (DE).			

(54) Title: A METHOD FOR STIMULATING THE IMMUNE SYSTEM



(57) Abstract

Medicament comprising a combination of at least one inhibitor of the effect of a substance negatively effecting an immune response, the substance selected from the group consisting of TGF-β and its receptors, VEGF and its receptors, interleukin 10 (IL-10) and its receptors, PGE₂ and its receptors, wherein the inhibitor has a molecular weight of less than 100 kDa and at least one stimulator positively effecting an immune response.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 99/04013

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 A61K48/00 A61K31/70 A61K38/20 A61K38/19 C12N15/11 //(A61K38/20,31:70),(A61K38/19,31:70)		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 A61P A61K C12N		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 02143 A (SIDNEY KIMMEL CANCER CENTER) 1 February 1996 (1996-02-01) page 4, line 16 -page 4, line 32 page 11, line 30 -page 12, line 1 page 12, line 22 -page 12, line 28 page 16, line 29 -page 17, line 9 example II	1-13
X	WO 94 25588 A (BIOGNOSTIK GES ;SCHLINGENSIEPEN GEORG FERDINAN (DE); BRYSCH WOLFGA) 10 November 1994 (1994-11-10) claim 1	10, 11
-/--		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents: <div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search <div style="text-align: center; font-size: 1.2em;">3 May 2000</div>		Date of mailing of the international search report <div style="text-align: center; font-size: 1.2em;">17. 05.00</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer <div style="text-align: center; font-size: 1.2em;">Pilling, S</div>

Form PCT/ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/04013

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JACHIMCZAK P ET AL: "The effect of transforming growth factor-beta 2-specific phosphorothioate-anti-sense oligodeoxynucleotides in reversing cellular immunosuppression in malignant glioma." JOURNAL OF NEUROSURGERY, vol. 78, no. 6, June 1993 (1993-06), pages 944-951, XP000886109 final paragraph of the "Materials and methods" on page 946 "discussion" on page 948	10,11
X	SPEARMAN M ET AL: "Antisense oligodeoxyribonucleotide inhibition of TGF - beta 1 gene expression and alterations in the growth and malignant properties of mouse fibrosarcoma cells." GENE, vol. 149, no. 1, 4 November 1994 (1994-11-04), pages 25-9, XP002089439 abstract; table 1	10,11
X	WO 95 00103 A (CHUNG HUN TAEG ;IL YANG PHARM CO LTD (KR)) 5 January 1995 (1995-01-05) claim 5	10,11
X	AKPORIAYE ET AL: "Concomitant expression of interferon-gamma and antisense TGF-beta transgenes enhances the immunogenicity of a murine mammary carcinoma" CANCER GENE THERAPY vol. 4, no. 6 conf. suppl. S55 6th International Conference on Gene Therapy of Cancer, San Diego, California, USA, Nov 20-22, 1997 XP000886176 abstract	1-13

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 99/04013

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 1-9, 12-13 (part)
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-9,12-13 (part)

Present Claim 1 relates to a medicament comprising a combination of;

i) an inhibitor of TGF-beta or TGF-beta receptors, VEGF or VEGF receptors, interleukin-10 or interleukin-10 receptors, PGE-2 or PGE-2 receptors and,

ii) a stimulator positively effecting an immune response

Each of these broad functionally defined categories of compounds (i) and (ii) includes many thousands of possible medicaments with the result that present Claim 1 comprises many, possible permutations. Furthermore, the Applicant is advised that, in general, compounds cannot be sufficiently defined by their function to the extent that a full search can be carried out.

Moreover, the definition of the "stimulator positively affecting the immune response" is vague and unclear.

Hence, in view of the above considerations a search of the full breadth of the present Claims 1 to 9 and 12 to 13 is impossible. Consequently, the search has been directed, in so far as practically feasible, to the general idea underlying the present invention and those combinations of medicaments which are most clearly described and exemplified namely combinations of;

i) an antisense oligonucleotide directed against TGF-beta or TGF-beta receptors, VEGF or VEGF receptors, interleukin-10 or interleukin receptors, PGE-2 or PGE-2 receptors, particularly the oligonucleotides of Claim 10 and;

ii) a stimulator selected from GM-CSF, SCF, CSF, IFN-gamma, FLT-3, MCP-1, IL-2, IL-4, IL-12, IL-18 (see page 3 lines 3 to 6); the stimulators described in categories (a), (b), (c) on present page 3; ras protein, p53 protein, EGF-receptor protein, retinoblastoma protein and derivatives thereof (see category (1) on present page 4).

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1 to 9, 12 and 13

Medicament comprising

- i) an inhibitor of TGF-beta or TGF-beta receptors, VEGF or VEGF receptors, interleukin-10 or interleukin receptors, PGE-2 or PGE-2 receptors in combination with;
- ii) a stimulator positively effecting an immune response

2. Claims: 10,11 (part)

An oligonucleotide having the sequence No. 55

3. Claims: 10,11 (part)

An oligonucleotide having the sequence No. 56

Each of the further sequences 57 to 213 relates to a further separate invention. Hence, inventions 4 to 158 are as defined herein above for invention 3. wherein oligonucleotide sequence No. 56 is replaced in turn by each of sequence No.'s 57 to 213.

i.e the present application comprises 160 inventions in total

With reference to the above identified inventions, it is noted that antisense oligonucleotides against TGF-beta at least are known (see each of the documents cited in the accompanying search report). Hence, there is no novel common inventive concept or idea linking the production of further anti-sense oligonucleotides as defined in Claims 10 and 11 which are directed against diverse targets such as TGF-beta, VEGF, flt, flkl and MCP-1.

Since all searchable claims could be searched without excessive additional effort justifying an additional fee, the present search report relates to ALL INVENTIONS IDENTIFIED ABOVE.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. National Application No

PCT/EP 99/04013

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9602143 A	01-02-1996	US 5772995 A	30-06-1998
		AU 711730 B	21-10-1999
		AU 3217895 A	16-02-1996
		CA 2195334 A	01-02-1996
		EP 0771147 A	07-05-1997
WO 9425588 A	10-11-1994	AU 6794594 A	21-11-1994
		EP 0695354 A	07-02-1996
		JP 8509370 T	08-10-1996
WO 9500103 A	05-01-1995	KR 9705347 B	15-04-1997
		AU 6984594 A	17-01-1995
		EP 0737071 A	16-10-1996
		JP 2548507 B	30-10-1996
		JP 7099977 A	18-04-1995
		US 5683988 A	04-11-1997
		ZA 9404185 A	08-02-1995

Form PCT/ISA/210 (patent family annex) (July 1992)